

N 66-16 382

FACILITY FORM 602

(ACCESSION NUMBER)
17
 (PAGES)
OK 69373
 (NASA CR OR TMX OR AD NUMBER)

(THRU)
1
 (CODE)
59
 (CATEGORY)

GPO PRICE \$ _____

CFSTI PRICE(S) \$ _____

Hard copy (HC) 1.00

Microfiche (MF) .50

ff 653 July 65

Department of Physics and Astronomy
THE UNIVERSITY OF IOWA

Iowa City, Iowa

A Correlation of Outer Radiation Zone
Electrons ($E_e \sim 1$ MeV) with the
Solar Activity Cycle*

by

L. A. Frank and J. A. Van Allen

Department of Physics and Astronomy
University of Iowa
Iowa City, Iowa

January 1966

*Research supported in part by the National Aeronautics and Space Administration under Grant NsG-233-62 and by the Office of Naval Research under Contract Nonr-1509(06).

Abstract

16382

A comparison of observations with several earth satellites reveals that the inner edge of the outer radiation zone of electron $E_e > 1.6$ MeV intensities has moved outward from $L \simeq 2.2$ in mid 1958 to $L = 3.0$ in late 1964. This outward motion of the outer radiation zone occurs during the declining phase of solar activity cycle 19 and strongly suggests a systematic change of the average value of one or more of the solar wind parameters over this time period.

Author

Introductory Comments

Measurements of the radiation belts surrounding the earth have now been performed over a substantial fraction of a solar activity cycle, the descending portion of solar cycle 19, with earth satellites and space probes beginning with the discovery of the geomagnetically trapped radiation with the instrumentation of Explorer 1 launched on 1 February 1958. The sun is now believed to be the predominant, primal source of energy required to maintain the radiation belts surrounding the earth--at least beyond an equatorial geocentric radial distance of about 2 earth radii--via the flow of solar corpuscular energy, or solar wind, past the earth. Thus it is of interest to combine the data from a number of satellites and space probes flown during the past seven years for an investigation of changes in the structure of the outer radiation zone during a solar activity cycle. One such study is reported here using an essentially homogeneous body of data obtained from similarly shielded ($\sim 1 \text{ g cm}^{-2}$), nominally identical Geiger-Mueller tubes which respond primarily to electrons $E_e \gtrsim 1 \text{ MeV}$ in the outer radiation zone [O'Brien, Van Allen, Laughlin, and Frank, 1962; Frank, Van Allen, and Hills, 1964], and which have been flown by us on Explorers 4, 7, 12, and 14; Injuns 1, 3, and 4; Pioneers 3 and 4; and OGO 1.

A Parameter for the Determination of the Long Term Variation

The profiles of electron ($E_e \gtrsim 1$ MeV) intensities in the outer radiation zone are characterized by marked temporal changes in gross structure and intensities in association with variations in geomagnetic activity as measured, for example, by the planetary magnetic indices Kp. Typically, during a geomagnetic storm cycle the intensities of electrons ($E_e \gtrsim 1$ MeV) fluctuate by a factor of ~ 100 and the position of the outer zone maximum intensity varies over a range of $L \simeq 3.0$ to 5.0 [Forbush, Pizzella, and Venkatesan, 1962] [Frank, Van Allen, and Hills, 1964]. Thus, in order to study variations of these profiles over a period of several years, it is necessary to find some characteristic of the signature of the outer radiation zone which is relatively insensitive to short time-scale (~ 1 month) perturbations. A feature of the intensity profiles utilized here is exemplified by the Explorer 12 data of Figure 1. Shown are three contours of the type 302 G.M. tube response for three similar Explorer 12 passes through the so-called 'slot', or minimum count-rate region at $L \simeq 2.6$ between the outer radiation zone at higher L-values and inner radiation zone (protons, $E_p \gtrsim 20$ MeV) at the lower L-values. Although these data were taken over a period of ~ 1 month during which the outer

zone maximum intensity changed markedly, the stability of the position of the inner edge of the outer radiation zone of electrons ($E_e \gtrsim 1$ MeV) at $L = 2.6 (+0.1)$ indicates that this parameter is appropriate for our present investigation of long term variations. Similar analyses of 302 G.M. tube data from Explorer 4 [Van Allen, McIlwain, and Ludwig, 1959], Pioneer 3 [Van Allen and Frank, 1959a], Explorer 7 [Forbush, Pizzella, and Venkatesan, 1962], Explorer 14 [Frank, Van Allen, and Hills, 1964], Injun 3 [Craven, 1966], and OGO 1 [Frank, Van Allen, Hills, and Fillius, 1965] are summarized in Figure 2. Note that data from both high and low eccentricity satellites and from space probes are combined in a single plot, and that determinations of the inner edge of the outer radiation zone as discussed above during similar periods for both high, eccentric and low, circular orbiting satellites are identical within experimental error (compare Pioneer 3 and Explorer 4, Explorer 14 and Injun 3). Electron ($E_e > 1.2$ MeV) measurements with a solid state, electron integral spectrometer on satellite 1963 38C [Williams and Kohl, 1965] have been included for comparison with the G.M. tube data. Figure 2 shows that the inner edge of the outer radiation zone has shifted from $L \simeq 2.2$ in mid 1958 to $L \simeq 3.0$ in late 1964, or $\Delta L \simeq 1$ over an approximately six-year period. The

gross nature of declining solar activity is represented by the parallel plot of smoothed sunspot numbers for this time period [Central Radio Propagation Laboratory Bulletin-F, Part B, issued January 1965] in Figure 2. The Explorer 14 and Injun 3 measurements shown in the summary graph may be artificially high due to the residual radiation injected by the three Soviet nuclear bursts during late October--early November 1962.

It may be inquired whether, for our present definition of the position of the inner edge of the outer radiation zone (Figure 1), the effect displayed in Figure 2 might be merely a manifestation of an increasing base-line 'slot' counting rate over the period of observations. Reference to the literature quoted above shows, however, that the 'slot' responses of the 302 G.M. tube at a given geomagnetic latitude are approximately constant or slowly decreasing over this time interval, thus producing a negligible, or slightly contrary effect.

Upon consideration of the above observational evidence we conclude that the inner edge of the outer radiation zone has shifted from $L \simeq 2.2$ to $L \simeq 3.0$ over the period 1958-1965 during the declining phase of the solar cycle.

Discussion

The most significant early study of the temporal variations of the electron distribution in the outer radiation zone is that of Forbush, Pizzella, and Venkatesan [1962]. These authors found a systematic negative correlation between the L value of the maximum outer zone intensity (L_{\max}) and the geomagnetic U-index during the period October 1959 to December 1960. The observed value of L_{\max} lay in the range 2.6 to 4.7, with a median value ~ 3.5 . During 1963, Armstrong [1965] found L_{\max} to have a median value ~ 4.0 with Injun 3; and during late 1964, Frank et al. [1965] found $L_{\max} \sim 4.5$ with OGO 1.

It is suggested that the outward movement of the inner edge of the outer zone by $\Delta L \sim 1.0$, as reported herein, and the outward movement of the peak intensity of the outer zone, also by $\Delta L \sim 1.0$, during approximately the same time period of declining solar activity are complementary consequences of the same basic cause--namely a systematic change of the average value of one or more of the solar wind parameters, density, bulk velocity, temperature, and magnetic field strength [cf. Snyder, Neugebauer, and Rao, 1963]. The physical linkage between the two measures of the outer zone profile is presumably the radial diffusion of ~ 1 MeV electrons as reported by Frank [1965], the average rate of diffusion

being apparently greater and/or the mechanism for radial diffusion penetrating more deeply into the magnetosphere during periods of higher average solar activity.

The solar-cycle variation of atmospheric density [King-Hele and Rees, 1965] is quantitatively inadequate [Van Allen, 1964] and of the wrong algebraic sign to account for the outward movement of the outer-zone profile.

Also it may be noted that the typical magnitude of the peak intensity of energetic electrons ($E_e \gtrsim 1$ MeV) in the outer zone near the geomagnetic equator as observed by Pioneers 3 and 4 [Van Allen and Frank, 1959a, b], Explorer 12 [Freeman, 1964], Explorer 14 [Frank, Van Allen, and Hills, 1964], and OGO 1 [Frank, Van Allen, Hills, and Fillius, 1965] has not displayed a marked change during the decreasing portion of the solar activity cycle, perhaps indicating a 'saturated' magnetosphere with regard to the population of ~ 1 MeV electrons.

In the framework of the present discussion, it is anticipated that the L-profile of the outer zone will move back inward toward the earth during the next several years of expected increase in solar activity.

REFERENCES

- Armstrong, T., Morphology of the outer zone electron distribution at low altitudes from January through July and September 1963 from Injun 3, J. Geophys. Res., 70, 2077-2110, 1965.
- Craven, J. D., The temporal variations of electrons at low altitudes in the outer radiation zone as observed with satellite Injun 3 (to be published, 1966).
- Forbush, S. E., G. Pizzella, and D. Venkatesan, The morphology and temporal variations of the Van Allen radiation belt, October 1959 to December 1960, J. Geophys. Res., 67, 3651-3668, 1962.
- Frank, L. A., Inward radial diffusion of electrons of greater than 1.6 million electron volts in the outer radiation zone, J. Geophys. Res., 70, 3533-3540, 1965.
- Frank, L. A., J. A. Van Allen, and H. K. Hills, A study of charged particles in the earth's outer radiation zone with Explorer 14, J. Geophys. Res., 69, 2171-2191, 1964.
- Frank, L. A., J. A. Van Allen, H. K. Hills, and R. W. Fillius, Proton and electron intensities in the earth's outer radiation zone: OGO 1 (title), Trans. Am. Geophys. Union, 46 (1), 124, 1965.

- Freeman, J. W., The morphology of the electron distribution in the outer radiation zone and near the magnetospheric boundary as observed by Explorer XII, J. Geophys. Res., 69, 1691-1723, 1964.
- King-Hele, D. G., and J. M. Rees, The decrease in upper-atmosphere density between 1957 and 1963, as revealed by satellite orbits, J. Atmos. Terrest. Phys., 25, 495-506, 1963.
- O'Brien, B. J., J. A. Van Allen, C. D. Laughlin, and L. A. Frank, Absolute electron intensities in the heart of the earth's outer radiation zone, J. Geophys. Res., 67, 397-403, 1962.
- Snyder, C. W., M. Neugebauer, and U. R. Rao, The solar wind velocity and its correlation with cosmic-ray variations and with solar and geomagnetic activity, J. Geophys. Res., 68, 6361-6370, 1963.
- Van Allen, J. A., Lifetimes of geomagnetically trapped electrons of several MeV energy, Nature (London), 203, 1006-1007, 1964.
- Van Allen, J. A., and L. A. Frank, Radiation around the earth to a radial distance of 107,400 kilometers, Nature (London), 183, 430-434, 1959a.

- Van Allen, J. A., and L. A. Frank, Radiation measurements to 658,300 km. with Pioneer IV, Nature (London), 184, 219-224, 1959b.
- Van Allen, J. A., C. E. McIlwain, and G. H. Ludwig, Radiation observations with satellite 1958 ϵ , J. Geophys. Res., 64, 271-286, 1959.
- Williams, D. J., and J. W. Kohl, Loss and replenishment of electrons at middle latitudes and high B values, J. Geophys. Res., 70, 4139-4150, 1965.

FIGURE CAPTIONS

Figure 1. Response of the 302 G.M. tube for several similar passes of Explorer 12 through the 'slot' region between the inner and outer radiation zones.

Figure 2. Summary of the positions of the inner edge of the outer radiation zone (electrons $E_e \gtrsim 1$ MeV) as observed with similar detectors in various spacecraft over a six year period. A measure of solar activity is given by the parallel plot of smoothed sunspot numbers.




Figure 1




FIGURE 2

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
University of Iowa, Department of Physics and Astronomy		UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE			
A Correlation of Outer Radiation Zone Electrons ($E_e \sim 1$ MeV) with the Solar Activity Cycle			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Progress			
5. AUTHOR(S) (Last name, first name, initial)			
Frank, L. A., and Van Allen, J. A.			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
January 1966		14	15
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
Nonr-1509(06)		U. of Iowa 66-1	
b. PROJECT NO.			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. AVAILABILITY/LIMITATION NOTICES			
Qualified requesters may obtain copies of this report from DDC.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		Office of Naval Research	
13. ABSTRACT			
<p>A comparison of observations with several earth satellites reveals that the inner edge of the outer radiation zone of electron $E_e > 1.6$ MeV intensities has moved outward from $L \simeq 2.2$ in mid 1958 to $L = 3.0$ in late 1964. This outward motion of the outer radiation zone occurs during the declining phase of solar activity cycle 19 and strongly suggests a systematic change of the average value of one or more of the solar wind parameters over this time period.</p>			

14.	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Outer Radiation Zone, Solar Wind						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.